



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

STATISTICAL REPORTS OF MORBIDITY AND MORTALITY, STATES
AND CITIES OF THE UNITED STATES (untabulated).

COLORADO.—Three months ended March 31, 1911. Population, 790,024. Total number of deaths from all causes 3,210, including diphtheria 24, measles 63, scarlet fever 44, tuberculosis 537, typhoid fever 40. Cases reported: Diphtheria 258, measles 2,316, scarlet fever 669, smallpox 757, typhoid fever 116.

FLORIDA.—Week ended June 24, 1911. Reports from the State board of health show malaria present in 2 localities (Jacksonville and Tampa) with 11 cases, smallpox in 8 counties with 27 cases, tuberculosis in 5 localities with 6 cases, typhoid fever in 1 locality (Tampa) with 8 cases.

INDIANA—*Terre Haute*.—Month of April, 1911. Population, 58,157. Total number of deaths from all causes 86. No deaths from contagious diseases. Cases reported: Diphtheria 7, measles 106, scarlet fever 20, smallpox 1, tuberculosis 1.

Four weeks ended May 27, 1911. Total number of deaths from all causes 74, including scarlet fever 1. Cases reported: Diphtheria 2, measles 49, scarlet fever 19, smallpox 1, tuberculosis 1, typhoid fever 2.

IOWA—*Ottumwa*.—Four weeks ended May 27, 1911. Total number of deaths from all causes 29, including scarlet fever 1, tuberculosis 2. Cases reported: Diphtheria 1, measles epidemic, scarlet fever 22, smallpox 1, tuberculosis 2.

KENTUCKY—*Louisville*.—Month of May, 1911. Population, 223,928. Total number of deaths from all causes 339, including diphtheria 2, measles 6, tuberculosis, pulmonary 37, typhoid fever 1. Cases reported: Diphtheria 14, measles 10, scarlet fever 21, smallpox 4, tuberculosis, pulmonary 50, typhoid fever 5.

MINNESOTA.—Month of March, 1911. Population, 2,075,708. Total number of deaths from all causes 2,007, including diphtheria 33, measles 17, scarlet fever 31, smallpox 1, tuberculosis 243, typhoid fever 25.

PENNSYLVANIA.—Month of May, 1911. Reports from the State department of health show a total of 12,848 cases of communicable diseases distributed as follows: Anterior poliomyelitis 4, cerebrospinal meningitis 16, chickenpox 883, diphtheria 865, erysipelas 184, German measles 71, leprosy 1, malarial fever 4, measles 5,714, mumps 615, pneumonia 268, puerperal fever 9, rabies 1, scarlet fever 1,092, smallpox 37, tetanus 2, trachoma 7, tuberculosis 1,291, typhoid fever 484, whooping cough 1,300.